

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

A61K 9/48H4^{Li6}



Europäisches Patentamt
European Patent Office
Office européen des brevets

Publication number:

0 274 870
A2

EUROPEAN PATENT APPLICATION

Application number: 87310931.8

Int. Cl. 4: A61K 9/10, A61K 47/00

Date of filing: 11.12.87

The title of the invention has been amended
(Guidelines for Examination in the EPO, A-III,
7.3).

Claims for the following Contracting States: ES
+ GR.

Priority: 18.12.86 GB 8630273

Date of publication of application:
20.07.88 Bulletin 88/29

Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

Applicant: T.I.L. Medical Ltd.
The Old Blue School Lower Square
Isleworth Middlesex TW7 6RL(GB)

Inventor: Story, Michael John
Elm Cottage Greaves Lane
Threapwood Near Malpas Cheshire SY14
6AS(GB)

Inventor: Flynn, Michael John
Hunterscombe Dorking Road
Leatherhead Surrey KT22 8JT(GB)

Representative: Sheard, Andrew Gregory et al
Kilburn & Strode 30, John Street
London WC1N 2DD(GB)

Micelles containing a non-steroidal antiinflammatory compound.

Non-steroidal anti-inflammatory drugs (NSAIDs) including diclofenac, flufenamic acid, flurbiprofen, ibuprofen, indomethacin, ketoprofen, naproxen, phenylbutazone, piroxicam and sulindac are administered in micelles to alleviate their adverse effects on the gastrointestinal tract. The drugs are formulated with surfactants such as polyethoxylated nonionics to give micelle-forming compositions.

EP 0 274 870 A2

Pharmaceutical Delivery Systems

This invention relates to pharmaceutical compositions for use in the treatment of inflammatory arthropathy.

Inflammatory arthropathy is the general name for a collection of debilitating and painful diseases which are extremely common in many countries of the world. Their classification is somewhat difficult, but inflammatory arthropathy or rheumatic disease seem to be the most common generic terms. In this specification, the term "inflammatory arthropathy" is used as the preferred generic term, but is to be understood to include forms of the disease known to some practitioners as rheumatic disease.

Of the various forms of inflammatory arthropathy, osteoarthritis (or osteoarthrosis) on the one hand and rheumatoid arthritis on the other hand are the commonest. Some workers in the field prefer the term osteoarthritis to the term osteoarthrosis, although it has been suggested that there is a place for both words. It has been suggested that osteoarthritis is the most sensible way of labelling the presence of simple degenerative joint disease but osteoarthrosis separates the acute episodes of an inflammatory nature which occur in degenerative joint disease.

Osteoarthritis usually has an insidious onset of pain, stiffness and a reduced range of movement. It commonly affects one or only a small number of joints. Intermittent swelling due to an effusion or an inflammatory episode in the affected joint may appear and, later in the disease, a permanent increase in size or change of shape may result from bony enlargement. Joint laxity develops with locking and grating.

It is often the joints which have been used the most or previously effected by trauma or inflammatory processes that suffer greatest damage. Thus, the weight-bearing joints of the hips and knees, the lumbar spine and the thumb bases (first carpometacarpal joints) are common victims of the disease. The latter are particularly effected in those who have been manual workers or even keen knitters.

The essential features of rheumatoid arthritis are pain and swelling of several joints with morning stiffness continuing for at least a few weeks. Rheumatoid arthritis tends to affect the peripheral small joints symmetrically. Whereas the joints in osteoarthritis may be described as dry, in rheumatoid arthritis they are "juicy", often swollen, hot, tender and red. There may also be accompanying systemic symptoms of a general malaise, weight loss, anorexia, mild fever and, on investigation, the finding of a normochromic (or hypochromic) normocytic anaemia.

Other common causes of inflammatory arthropathy include viral arthritis, ankylosing spondylitis, psoriatic arthropathy, Reiter's disease, gouty arthritis, septic arthritis (suppurative arthritis), erythema nodosum and Henoch-Schoenlein purpura. The most important in the present context are ankylosing spondylitis and gouty arthritis.

Ankylosing spondylitis is characterised by the gradual onset of low-back pain (sometimes bilateral buttock pain) with morning stiffness. Peripheral joints may become effected. There is a reduced range of spinal movement and chest expansion. Rigidity of the spine follows, often in a cranial direction (first lumbar, then dorsal then cervical) with a characteristic clinical picture of high dorsal kyphosis, obliteration of lumbar lordosis and flattening of the chest.

Gouty arthritis is due to the deposition of monosodium urate monohydrate crystals in the joint. Gouty arthritis is a very common disease: it is estimated that there are over 300,000 sufferers in the United Kingdom alone. The popularly held belief that gout is largely due to an over indulgence of port and pheasant is mainly fallacious, although provocative factors may often be related to its onset. Examples include trauma, surgery, unusual physical exercise, severe illness, dietary excess, alcohol and drugs. Any joint may be affected, and the onset may be polyarticular. Affected joints are painful, red, hot, swollen and exquisitely tender.

The treatment of inflammatory arthropathy has naturally received a fairly large amount of attention from pharmacologists and pharmaceutical manufacturers. A first class of drugs that have been used in the treatment of inflammatory arthropathy are steroids. Cortisol and its synthetic analogues have the capacity to prevent or suppress the development of the local heat, redness, swelling and tenderness by which inflammation is recognised. At the microscopic level they inhibit not only the early phenomena of the inflammatory process (oedema, fibrin deposition, capillary dilation, migration of leukocytes into the inflamed areas and phagocytic activity) but also the later manifestations (capillary proliferation, fibroblast proliferation, deposition of collagen and, still later, cicatrization).

In clinical terms, the administration of such corticosteroids for their anti-inflammatory effects is palliative therapy. The underlying cause of the disease remains; the inflammatory manifestations are merely suppressed. Nevertheless, they are effective in affording symptomatic relief, but prolonged administration of corticosteroids may be a very high price to pay for such relief: the adrenal cortex may become atrophied,

thereby limiting the body's own ability to survive and adapt in a constantly changing environment. The adrenal cortex is the organ of homeostasis: in the absence of the adrenal cortex, survival is possible, but only under the most rigidly prescribed conditions. In more general terms, it has long been recognised that corticosteroids are powerful drugs with slow cumulative toxic effects on many tissues, which may not be apparent until made manifest by a catastrophe.

In the treatment of inflammatory arthropathy, the focus of attention shifted from steroids to a structurally unrelated group of compounds known as slow acting anti-rheumatic drugs (SAARDs). SAARDs have empirically been categorised into three groups. Group I, including drugs of proven value which are widely used, encompasses azathioprine, chloroquine, D-penicillamine and gold salts. Group II relating to clinically active drugs under continuing investigation, includes cyclophosphamide, dapsone, levamisole, methotrexate, sulphasalazine, thiols and thymopoietin. The group III SAARDs are those of less practical or unproven treatment: this group includes methylprednisolone pulsing.

The range of SAARDs is considerable, as has been seen above, and despite much experimental work their modes of action are largely unknown. Logistical and toxicity factors prevent the use of SAARDs in all patients.

A third category of drugs for use in the treatment of inflammatory arthropathy consists of the non-steroidal anti-inflammatory drugs (NSAIDs). Aspirin is the prototype NSAID, and for this reason this group of drugs is also known as the "aspirin-like" drugs. This secondary nomenclature gives a key to a functional similarity of NSAIDs in the absence of any overall chemical similarity: they all appear to owe their anti-inflammatory action, at least in part, to the inhibition of prostaglandin synthesis. According to Goodman and Gilman in "The Pharmacological Basis of Therapeutics" MacMillan 7th Edition 1985, it has been established in recent years that:

1. All mammalian cell types studied (with the exception of the erythrocyte) have microsomal enzymes for the synthesis of prostaglandins;

2. Prostaglandins are always released when cells are damaged and have been detected in increased concentrations in inflammatory exudates - all available evidence indicates that cells do not store prostaglandins, and their release thus depends on biosynthesis *de novo*;

3. All aspirin-like drugs inhibit the biosynthesis and release of prostaglandins in all cells tested; and

4. With the exception of the anti-inflammatory glyocorticoids, other classes of drugs generally do not affect the biosynthesis of prostaglandins.

NSAIDs (or aspirin-like drugs - the two terms are used interchangeably in this specification) can be categorised conveniently into six structural groups. First, there are the salicylic acids and esters including aspirin, benorylate, aloxiprin, salsalate and choline magnesium trisalicylate.

Secondly, there are the propionic acid derivatives, including ibuprofen, naproxen, flurbiprofen, ketoprofen, fenoprofen, fenbufen, benoxaprofen and suprofen.

Thirdly, there is the class of oxicams, including piroxicam.

Fourthly, acetic acid derivatives can be split into two subclasses. Phenylacetic acids include diclofenac and fenclofenac; carbo- and heterocyclic acetic acids include indoles such as indomethacin and sulindac and pyrroles such as tolmetin.

Fifthly, there are the pyrazolones which include oxyphenbutazone, phenylbutazone, feprazone and azapropazone.

Sixthly, the fenamic acid derivatives include flufenamic acid and mefenamic acid.

NSAIDs have emerged as the drugs of choice in the treatment of inflammatory arthropathy. This is possibly more due to the disadvantages associated with other classes of drugs than in anything else. As indicated previously, the inflammatory diseases of the joints cause an extremely high level of discomfort and in many instances the results are crippling. The requirement for treatment is unquestioned and the treatment is in many cases chronic, that is to say it is continuous as the diseases are generally incurable. Unfortunately, the common element in the therapeutic properties of the NSAIDs is also the principle cause of side effects. As has been mentioned, the salicylates and other NSAIDs are thought to be effective in inflammatory joint disease, and their effectiveness is thought to be partly mediated through prostaglandin inhibition. Prostaglandins have been shown to have a protective effect on the gastrointestinal mucosa and, therefore, drugs which inhibit their activity are likely to cause gastrointestinal intolerance. Drugs with a potent inhibitory action on prostaglandin synthetase are marketed as having a potent anti-inflammatory action but have been shown to cause more faecal blood loss than those with weak anti-prostaglandin activity. Aspirin, for example, causes as much as an 8-to 10-fold increase in faecal blood loss and indomethacin a nearly 3-fold loss, compared with controls. However, when oral prostaglandin E2 (PGE2) at doses of 1mg three or four times daily is given with indomethacin or aspirin, the blood loss is reduced to control levels without reducing the effectiveness of the drugs.

Protection of the stomach from the drug has in some circumstances been shown to be effectively achieved by the use of enteric coating, as demonstrated by enteric coated aspirin preparations. However, the use of conventional enteric coating means that the drug is released in the neutral or slightly alkaline environment of the small or large intestine, which consequently experiences a considerably heightened local concentration from direct contact by the drug. Intestinal ulceration can occur with chronic administration of NSAIDs.

There is therefore a need for an improved and safer form of administration of NSAIDs to give protection both in the stomach and in the intestine. In addition, it would be advantageous to be able to provide a means of enhancing the absorption of the NSAIDs, which tend to be poorly water soluble, as well as providing an improved concentration of the drug at the cellular level at the site of its action. It is known that drugs with a low water solubility have a slow and variable dissolution pattern which can lead to reduced and erratic bioavailability. In short, what has been needed for some time is a delivery system for NSAIDs which protects the gastrointestinal tract from the drug, and which provides a means of alleviating the difficulties associated with very poor water solubility.

The present invention is based on the discovery that the use of micelles enables a particularly appropriate form of administration of NSAIDs to be achieved.

According to a first aspect of the present invention, there are provided micelles containing a non-steroidal anti-inflammatory drug.

Although NSAIDs themselves tend not to form micelles, amphipathic compounds, known more familiarly as surfactants, can form micelles. Surfactants have two distinct regions in their chemical structure, termed hydrophilic (water-liking) and hydrophobic (water-hating) regions. Micelles are aggregates in which the surfactant molecules are generally arranged in a spheroidal structure with the hydrophobic region at the core shielded, in an aqueous solution, from the water by a mantle of outer hydrophilic regions. According to a second aspect of the invention, therefore, there is provided a pharmaceutical composition comprising a non-steroidal anti-inflammatory drug and a surfactant, the composition being capable of forming micelles containing the non-steroidal anti-inflammatory drug when administered orally. It will generally be the case that the drug will be dissolved in the surfactant. In its simplest form, the pharmaceutical composition can be a solution of the drug in a surfactant, although other components may be present in the system if desired or necessary.

In a third aspect, the invention provides a process for the preparation of an anti-inflammatory composition capable of forming non-steroidal anti-inflammatory drug-containing micelles on oral administration to a human or non-human animal, the process comprising admixing a non-steroidal anti-inflammatory drug with a surfactant. The process may involve dissolving the drug in the surfactant.

According to a fourth aspect, the invention provides the use of a non-steroidal anti-inflammatory drug and a surfactant in the preparation of a composition for administering the drug in micellar form. Insofar as the law allows, the invention also relates to a method for the treatment or prophylaxis of inflammatory arthropathy, the method comprising the administration of micelles containing a non-steroidal anti-inflammatory drug.

Micelles are to be contrasted in terms of their structure with vesicles and with liposomes. Vesicles are aggregates of amphipathic molecules arranged in a bilayer. Typically, a vesicle will have a hydrophilic interior and a hydrophilic exterior: hydrophilic regions of an internal layer of the molecules will be directed inwardly, and hydrophilic regions of an outer layer of the molecule will be directed outwardly. Hydrophobic regions of the two layers will be directed towards one another within the molecular wall of the vesicle.

Liposomes are nothing more than multilamellar vesicles, as is revealed by the fact that liposomes disintegrate to vesicles upon ultrasonication.

Surfactants can be variously classified, and often by reference to the nature of the hydrophilic region, which can be anionic, cationic, zwitterionic or non-ionic. In the present invention, nonionic surfactants are preferred. A particularly preferred subcategory of nonionic surfactants are polyoxyethylated surfactants, including polyoxyethylated glycol monoethers, polyoxyethylated fatty acids, polyoxyethylated sorbitan fatty esters, and polyoxyethylated castor oils. However, other nonionic surfactants are also particularly appropriate, including sorbitan fatty acid esters, poloxamers, polyethylene glycol fatty acid esters and polyethoxylated glyceryl fatty acid esters.

Whatever the precise chemical structure of the surfactant or surfactants used, it is generally preferred to use one or more of those that have been already cleared for human ingestion. Therefore, surfactants with a low toxicity are preferred. For example, surfactants having an LD₅₀ exceeding 10 g/kg and preferably 15 g/kg, are generally suitable. The absence of other side effects is of course also appropriate. Although surfactants which have already been approved for human ingestion are naturally preferred, the use of other

surfactants is not ruled out, not least because they may in time come to be approved for human ingestion. The availability of nonionic surfactants is not perceived to be a cause of difficulty. For example, the following surfactants are known to be available.

5 Polyoxyethylene Alkylphenols POE(n) octylphenol n = 1-70
 Triton X series (Rohm & Haas) Igepal CA series (GAF, USA) Antarox CA series (GAF, UK)
 POE(n) nonylphenol n = 1.5-100
 Triton N series (Rohm & Haas) Igepal CO series (GAF, USA) Antarox CO series (GAF, UK)

10 None of the polyoxyethylene alkylphenols are as yet approved for human ingestion.

15 Polyoxyethylated Glycol Monoethers POE(n) lauryl ether n = 4,23
 Volpo L series (Croda)
 Brij 30 series (Atlas ICI Specialties, UK)
 POE(n) cetyl ether n = 2,10,20
 Brij 50 series (Atlas ICI)
 20 POE(n) stearyl ether n = 2,10,20
 Brij 70 and 700 series (Atlas ICI)
 POE(n) oleyl ether n = 2-20
 Volpo N series (Croda)
 Brij 90 series (Atlas ICI)
 25 POE(n) ceto-stearyl ether n = 3-20
 Volpo CS series (Croda)

30 None of these have been approved for internal use, although Cetomacrogol 1000 (Brij 58, Volpo CS20) has been extensively used in topical applications.

Polyoxyethylated Glyceryl Fatty Acid Esters

35 POE(n) glyceryl monolaurate n = 15,40 Glycerox L series (Croda)
 These products have not been cleared for internal ingestion.

40 Polyoxyethylated Fatty Acids POE(n) monolaurate n = 4-100
 Crodet L series (Croda)
 POE(n) monooleate n = 4-100
 Crodet O series (Croda)
 POE(n) monostearate n = 4-100
 Crodet S series (Croda)
 45 Myrij series (Atlas ICI)

50 POE(8) monostearate and POE(40) monostearate appear to be approved for internal ingestion in the UK and EEC, and the latter is also approved by the FDA in the US. The other POE(n) monostearates appear valid contenders for approval, with the POE(n) monooleates and monolaurates also being likely candidates.

55 Sorbitan Fatty Acid Esters Sorbitan monolaurate
 Crill 1 (Croda)
 Span 20 (Atlas ICI)
 Sorbitan monopalmitate
 Crill 2 (Croda)
 Span 40 (Atlas ICI)

Sorbitan monostearate
 Crill 3 (Croda)
 Span 60 (Atlas ICI)
 Sorbitan tristearate
 5 Crill 35 (Croda)
 Span 65 (Atlas ICI)
 Sorbitan monooleate
 Crill 4 (Croda)
 Span 80 (Atlas ICI)
 10 Sorbitan sesquioleate
 Crill 43 (Croda)
 Sorbitan trioleate
 Crill 45 (Croda)
 Span 85 (Atlas ICI)
 15 Sorbitan monoisostearate
 Crill 6 (Croda)

20 The surfactants in this group have good approval rating in the UK, EEC and US, but not complete approval.

Polyoxyethylated Sorbitan Fatty Acid Esters POE(20) sorbitan monolaurate
 Crillet 1 (Croda)
 25 Tween 20 (Atlas ICI)
 POE(4) sorbitan monolaurate
 Crillet 11 (Croda)
 Tween 21 (Atlas ICI)
 POE(20) sorbitan monopalmitate
 30 Crillet 2 (Croda)
 Tween 40 (Atlas ICI)
 POE(20) sorbitan monostearate
 Crillet 3 (Croda)
 Tween 60 (Atlas ICI)
 35 POE(4) sorbitan monostearate
 Crillet 31 (Croda)
 Tween 61 (Atlas ICI)
 POE(20) sorbitan tristearate
 Crillet 35 (Croda)
 40 Tween 65 (Atlas ICI)
 POE(20) sorbitan monooleate
 Crillet 4 (Croda)
 Tween 80 (Atlas ICI)
 POE(5) sorbitan monooleate
 45 Crillet 41 (Croda)
 Tween 81 (Atlas ICI)
 POE(20) sorbitan trioleate
 Crillet 45 (Croda)
 Tween 85 (Atlas ICI)
 50 POE(20) sorbitan monoisostearate
 Crillet 6 (Croda)

55 These surfactants have a similar approval profile to the Sorbitan Fatty Acid Esters, above.

Polyoxyethylated Castor Oils POE(n) castor oil n = 10-100

Etocas Series (Croda)

Cremophor EL (BASF)

POE(n) hydrogenated castor oil n = 10-100

5 Croduret series (Croda)

Cremophor RH40 (BASF)

10 Cremophor EL and Cremophor RH40 are well established as orally ingestible surfactants. It is envisaged that there would be no problems in registering the Etocas or Corduret series provided BP Castor Oil was used in manufacture of the surfactant.

Poloxamers POE(n)-POP(m)

15 Syneronic PE series (ICI Petrochem & Plastics Div) Pluronic series (Wyandotte Chem. Corp. USA)

Some of these have been used in orally ingested pharmaceuticals. They are of low toxicity.

20 Polyethylene Glycol Fatty Acid Esters PEG(400) distearate

Cithrol 4DS (Croda)

PEG(400) monolaurate

Cithrol 4ML (Croda)

25 PEG(n) monooleate n = 200,300,400

Cithrol MO series (Croda)

PEG(400) dioleate

Cithrol 4DO (Croda)

PEG(n) monostearate n = 400,600 1000

30 Cithrol MS series (Croda)

There are no toxicology data readily available for these surfactants.

One factor affecting the choice of surfactant or surfactants to be used is the hydrophilic-lipophilic balance (HLB), which gives a numerical indication of the relative affinity of the surfactant for aqueous and non aqueous systems. Surfactants having an HLB of about 10 or above, particularly about 12 or above, are preferred. However, there may be cases where a mixture of two or more surfactants provides an improved degree of solubilization over either surfactant used alone.

In addition to the HLB, the nature of the hydrophobic chain may be taken into account. For example, increasing the degree of unsaturation may improve the potential for solubilization, as may increasing the chain length and/or having branches. Further a reduction in the molecular weight may give improved solubilization on a weight for weight basis, even at the expense of a slight reduction in the HLB. It has been discovered that it is the provision of the solubilizing interior of the micelles which is important, and this may be related to the formation of a solution of the drug in the surfactant prior to the addition of the aqueous phase.

45 The physical nature of the surfactants will also be a factor to be taken into consideration when choosing surfactants for a particular formulation. The choice of surfactant will, among other things, depend on the type of formulation. For example, a formulation in the form of a solution may be in the form of a liquid, although a solid surfactant may be used in formulating a solution. Soft gelatin capsules may be formulated using a surfactant in the form of a liquid, a viscous liquid or melted waxy solid. Hard gelatin capsules may be formulated using a liquid, a paste (melted) or a solid (melted) surfactant. There follows below a list of potential nonionic surfactants, together with a description of their physical nature and an indication of their HLB and LD₅₀.

Chemical Identity	Description	HLB	LD50 g/kg
<u>Polyoxyethylated Glycol Monoethers</u>			
POE(4) lauryl ether	Water white liquid	9.5	9
POE(23) lauryl ether	Off-white soft solid	17.0	9
POE(2) cetyl ether	White solid	5.3	22
POE(15) cetostearyl ether	Off-white waxy solid	14.6	?
POE(20) cetostearyl ether	Off-white hard waxy solid	15.6	3.6
POE(15) oleyl ether	Pale straw paste	14.2	?
POE(20) oleyl ether	Pale straw soft solid	15.5	15.1
POE(2) stearyl ether	White solid	4.9	>25
POE(2) oleyl ether	Pale yellow liquid	4.9	25
<u>Polyoxyethylated Fatty Acids</u>			
POE(4) monolaurate	Pale straw liquid	9.3	?
POE(8) monolaurate	White soft solid	12.7	?
POE(12) monolaurate	White soft solid	14.5	?
POE(24) monolaurate	White waxy solid	16.8	?
POE(40) monolaurate	White hard solid	17.9	?
POE(100) monolaurate	White hard solid	19.1	?
POE(4) monooleate	Yellow/amber liquid	7.7	?
POE(8) monooleate	Yellow/amber liquid	10.4	?
POE(12) monooleate	Yellow/amber liquid	13.4	?
POE(24) monooleate	Yellow/amber paste/solid	15.8	?
POE(40) monooleate	Yellow soft solid	17.4	?

Chemical Identity	Description	HLB	LD50 g/kg
POE(100) monooleate	Yellow waxy solid	18.8	?
POE(4) monostearate	White soft waxy solid	7.7	?
POE(8) monostearate	White waxy solid	11.1	64
POE(12) monostearate	White waxy solid	13.4	?
POE(20) monostearate	White waxy solid	15.0	10
POE(24) monostearate	White waxy solid	15.8	?
POE(30) monostearate	White hard solid	16.0	?
POE(40) monostearate	White hard solid	16.9	>30
POE(50) monostearate	White hard solid	17.9	>25
POE(100) monostearate	White hard solid	18.8	25
<u>Sorbitan Fatty Acid Esters</u>			
Sorbitan monolaurate	Pale yellow viscous liquid	8.6	41
Sorbitan monopalmitate	Pale tan waxy solid	6.7	>16
Sorbitan monostearate	Pale tan waxy solid	4.7	31
Sorbitan tristearate	Pale tan waxy solid	2.1	>16
Sorbitan monooleate	Amber viscous liquid	4.3	>40
Sorbitan sesquioleate	Amber viscous liquid	3.7	?
Sorbitan trioleate	Amber viscous liquid	1.8	>40
Sorbitan monoisostearate	Yellow viscous liquid	4.7	?

Chemical Identity	Description	HLB	LD50 g/kg
<u>Polyoxyethylated Sorbitan Fatty Esters</u>			
POE(20) sorbitan monolaurate	Pale yellow liquid	16.7	>39
POE(4) sorbitan monolaurate	Yellow/amber liquid	13.3	>38
POE(20) sorbitan monopalmitate	Yellow pasty liquid	15.6	>38
POE(20) sorbitan monostearate	Yellow pasty liquid	14.9	>38
POE(4) sorbitan monostearate	Pale yellow waxy solid	9.6	>40
POE(20) sorbitan tristearate	Cream waxy solid	10.5	>40
POE(20) sorbitan monooleate	Yellow/amber liquid	15.0	>38
POE(5) sorbitan monooleate	Yellow/amber liquid	10.0	>37
POE(20) sorbitan trioleate	Yellow/amber liquid	11.0	>36
POE(20) sorbitan monoisostearate	Yellow liquid	14.9	?
<u>Polyoxyethylated Castor Oils</u>			
POE(10) castor oil	Pale yellow liquid	6.3	?
POE(35) castor oil	Pale yellow liquid	12.5	>10
POE(40) castor oil	Pale yellow liquid	13.0	?
POE(60) castor oil	Pale yellow soft paste	14.7	?
POE(100) castor oil	Pale yellow waxy solid	16.5	?
POE(10) hydrogenated castor oil	Pale straw liquid	6.3	?
POE(30) hydrogenated castor oil	Pale straw liquid	11.6	?
POE(40) hydrogenated castor oil	White soft paste	13.0	?
POE(45) hydrogenated castor oil	White soft paste	14	>16

Chemical Identity	Description	HLB	LD50 g/kg
POE(60) hydrogenated castor oil	White soft paste	14.6	?
POE(100) hydrogenated castor oil	White waxy solid	16.4	?
<u>Poloxamers</u>			
POE(22) - POP (13)	Liquid	18.5	
POE(90) - POP (13)	Solid	30.5	
POE(7) - POP (17)	Liquid	8	
POE(20) - POP (17)	Liquid	16	
POE(4) - POP (23)	Liquid	3	
POE(10) - POP (23)	Liquid	7	
POE(27) - POP (23)	Liquid	7	
POE(159) - POP (23)	Solid	15	
POE(47) - POP (27)	Paste	16.5	
POE(6) - POP (30)	Liquid	2	
POE(51) - POP (30)	Paste	16	
POE(119) - POP (30)	Solid	24	
POE(205) - POP (30)	Solid	28	
POE(19) - POP (37)	Liquid	5.5	
POE(41) - POP (37)	Paste	13.5	
POE(8) - POP (43)	Liquid	1	
POE(32) - POP (43)	Paste	9	
POE(296) - POP (43)	Solid	27	
POE(10) - POP (53)	Liquid	0.5	
POE(193) - POP (53)	Solid	22	

Various non-steroidal anti-inflammatory drugs in common use today tend to have, as a common property, the property of being poorly soluble in water. The poor solubility does nothing to ameliorate the problems of their administration in conventional delivery systems, and the present invention provides a means of overcoming at least some of the difficulties associated with poor water solubility. Apart from anything else, particles of insoluble drug may tend to lie in folds of the intestinal mucosa, thereby giving rise to local irritancy.

There follows a brief discussion of each of the NSAIDs which are, in accordance with the present invention, particularly appropriate for being delivered in the form of micelles.

Diclofenac is sold as the free acid under the trade mark VOLTAROL by Geigy Pharmaceuticals. It is poorly soluble in water but soluble in some organic solvents. Gastrointestinal disturbances have been reported in about 7% of all cases. In general, it is fairly well absorbed, but more than 99% of the drug has been found to be bound to plasma proteins. The drug has been recommended for use in the treatment of rheumatoid arthritis and other rheumatic disorders at a dose of from 75 to 150 mg per day, depending upon the form of administration and its frequency. Diclofenac has been supplied as enteric coated tablets, slow release tablets, suppositories and in ampoules.

Flufenamic acid is sold under the trade mark MERALEN by Merrell Dow Pharmaceuticals. Its solubility is less than 1 part in 10,000 parts of water, although it is reasonably soluble in various organic solvents. Its most frequent adverse effects are gastrointestinal disturbances. The drug is well absorbed and is extensively bound to plasma proteins. It is prescribed for rheumatic disorders at doses of from 400 to 600 mg per day.

Flurbiprofen is sold under the trade mark FROBEN by the Boots Company plc. It is soluble in 100 to 1,000 parts of water only, but is readily soluble in most organic solvents. Gastrointestinal side effects have been reported in from 23 to 27% of cases. It is readily absorbed, approximately 99% of the drug being bound to plasma proteins. It is prescribed for rheumatoid arthritis and other rheumatic disorders and doses from 150 to 200 mg per day in a divided dose. The maximum dosage is stated to be 300 mg per day.

Another Boots Company drug is ibuprofen sold under the trade mark BRUFEN. Other trade marks in the UK for ibuprofen are FENBID and APSIFEN and in the US are RUFEN, ADVIL, MOTRIN and NUPRIN. It is poorly soluble in water: less than 1 part of drug will dissolve in 10,000 parts of water. However, it is fairly soluble in simple organic solvents. The most frequent adverse effects reported are, again, gastrointestinal. The drug is well absorbed and extensively bound to plasma proteins in vivo. It is prescribed for rheumatic arthritis and other musculoskeletal disorders, as well as acute gout. The dosage of the drug is from 600 to 1200 mg daily in divided doses, with 2,400 mg per day being the maximum.

Indomethacin is sold under the trade mark INDOCID by Thomas Morson Pharmaceuticals. It is also sold under the trade mark INBRILON in the UK and INDOCIN in the US. One part of drug is only soluble in more than 10,000 parts of water, but is more soluble in simple organic solvents. The most frequently reported adverse effects are gastrointestinal problems, headache and dizziness. The drug is readily absorbed, with more than 90% being bound to plasma proteins. It is prescribed for rheumatoid arthritis, ankylosing spondylitis, osteoarthritis and other rheumatic disorders, as well as acute gout. The recommended dosage is up to 150 to 200 mg daily in divided doses.

Ketoprofen is sold under the trade mark ORUDIS by May & Baker Limited, who also market controlled release pellets of the drug under the trade mark ORUVAIL. It is also sold in the UK under the trade mark ALRHEUMAT. Its solubility is less than 1 part in 10,000 parts of water, but it is freely soluble in various simple organic solvents. The most frequent side effects are gastrointestinal. The drug is readily absorbed and is extensively bound to plasma proteins. It is prescribed for rheumatoid arthritis and osteoarthritis at doses of from 50 to 100 mg twice daily.

Naproxen is sold under the trade mark NAPROSYN by Syntex Pharmaceuticals Limited. Naproxen sodium is sold as SYNFLEX. The solubility of the free acid is less than 1 part in 10,000 parts water, but the drug is more soluble in simple organic solvents. The most frequent adverse effects reported are gastrointestinal. The drug is readily absorbed with more than 99% being bound to plasma proteins. Naproxen is prescribed for rheumatoid arthritis and other rheumatic or musculoskeletal disorders, dysmenorrhoea and acute gout. Its recommended dosage is from 500 to 1,000 mg daily in divided doses, with from 250 to 375 mg twice daily being preferred.

Phenylbutazone has been sold in the UK under the trade mark BUTAZOLIDIN by Geigy Pharmaceuticals; it is still available in the United States. Its solubility is less than 1 part in 10,000 parts of water, but it is more in common organic solvents. Its most adverse effects are nausea, vomiting and epigastric distress. It is readily absorbed, with 98% of the drug being bound to plasma proteins. It is generally only prescribed for the treatment of rheumatic disorders where other drugs have failed. The initial recommended

dosage ranges from 400 to 600 mg per day, but this should decrease to a maintenance dosage of from 200 to 300 mg per day. In both cases, the dosages should be divided through the day. The maximum daily dosage is 800 mg.

Piroxicam is marketed in the UK under the trade mark FELDENE by Pfizer Limited. It is known to be poorly soluble in water but soluble in some organic solvents. There is a high incidence of severe gastrointestinal side-effects. The drug is well absorbed with 99% being bound to plasma proteins. It is prescribed for rheumatoid arthritis and other rheumatic disorders, as well as acute gout at dosages of from 10 to 30 mg per day, with 20 mg per day being preferred.

Sulindac is sold in the UK under the trade mark CLINORIL by Merck, Sharp & Dohme Limited. Its solubility is less than 1 part in 10,000 parts water, although it is slightly soluble in simple organic solvents. The most frequent side effects claimed of are gastrointestinal, headache and dizziness. It is incompletely absorbed from the gastrointestinal tract. It is prescribed for rheumatic and other musculoskeletal disorders at dosages of from 400 to 600 mg per day.

Specific paediatric preparations include:

Ibuprofen 200 ml \times 100 mg:5 ml syrup;

Indomethacin 200 ml \times 25 mg:5 ml suspension (UK, but not recommended in US for children under 14 years); and

Naproxen 500 ml \times 25 mg/ml suspension.

Ketoprofen appears to be a possible further candidate for paediatric use.

Various surfactants and NSAIDs suitable for use in the present invention have now been described. However, the list is not to be taken as exhaustive. In addition, it should not be assumed that only these two ingredients have to be present as in some cases, including capsules, anti-oxidants will be required to ensure adequate stability. When preparing solutions, for example, for paediatric or geriatric use, additional excipients may be present such as preservatives, sweeteners and flavouring agents.

In certain cases it may be required to formulate an NSAID capsule which has sustained release properties. In such cases it is appropriate to include in the formulation ingredients which slow down the release of the surfactant NSAID combination from the total capsule mix. Such ingredients will generally be of a waxy nature, but this will not exclude the opportunity of using other techniques such as pellets with controlled release coatings.

The relative proportions of drug and surfactant used will, in the main, depend upon (a) the drug, (b) the surfactant and (c) the intended formulation, be it hard gelatin capsules, liquid solution or whatever. When preparing a micelle-forming drug/surfactant mix for use in capsules, it may be found appropriate to use the drug and surfactants in a weight ratio (drug:surfactant) of from 1:5.7 to 1:50, for example, from 1:6 to 1:20 or 1:25. When preparing solutions for, for example, paediatric or geriatric use, the drug:surfactant ratio may range from 1:8 to 1:30, with from 1:10 to 1:27.5 being preferred.

The following examples illustrate the invention.

EXAMPLE 1

Indomethacin Capsules - Size 2

Capsules of 25 mg active ingredient per capsule were prepared using the following proportions:

	<u>mg per capsule</u>
Indomethacin	25
POE(20) sorbitan monooleate (CRILLET 4)	310
Total	335

The surfactant is heated to 50-60°C and the active ingredient is then added with stirring, the latter being sufficiently vigorous to ensure that the active ingredient dissolves completely in the surfactant.

When the mixture is homogeneous and it becomes a clear solution, it is stirred for at least a further 15 minutes before filling into capsules, the temperature being maintained at 50-60°C.

The filling of capsules requires equipment the same or similar to that used for filling Licaps of Capsugel. The capsule used in this example is the Licaps hard gelatin capsule, size 2. The capsule is filled to approximately 90% of its nominal capacity to ensure that there is no spillage, and the cap is sealed onto the body by the Licaps sealing process. This ensures no leakage of liquid contents, or of solid contents which may melt if raised to a moderately high temperature during transport, as well as providing security against tampering.

EXAMPLES 2 TO 11

The procedure of Example 1 was repeated except that 310 mg capsule of the surfactant indicated below was used.

In all cases the drug:surfactant weight ratio was 1:12.4.

<u>Example No</u>	<u>Surfactant</u>
2	POE(20) sorbitan monoisostearate (CRILLET 6)
3	POE(40) monostearate (CRODET S24)
4	POE(24) monostearate (CRODET S40)
5	POE(40) monooleate (CRODET O40)
6	POE(20) cetostearyl ether (VOLPO CS20)
7	POE(15) cetostearyl ether (VOLPO CS15)
8	POE(20) oleyl ether (VOLPO N20)
9	POE(15) oleyl ether (VOLPO N15)
10	POE(40) hydrogenated castor oil (CREMOPHOR RH40)
11	POE(35) castor oil (ETOCAS 35)

EXAMPLE 12

Indomethacin Capsules - Size 1

Following the procedure of Example 1, but using Size 1 capsules, capsules of 25 mg active ingredient per capsule were prepared using the following proportions:

	<u>mg per capsule</u>
Indomethacin	25
POE(20) sorbitan monooleate (CRILLET 4)	425
Total	450

EXAMPLES 13 TO 23

The procedure of Example 12 was repeated except that 425 mg. capsule of the surfactant indicated below was used. In all cases the drug:surfactant weight ratio was 1:17.

5

<u>Example No</u>	<u>Surfactant</u>
10	13
	POE(20) sorbitan monoisostearate (CRILLET 6)
	14
	POE(40) monostearate (CRODET S40)
15	15
	POE(24) monostearate (CRODET S24)
	16
	POE(40) monooleate (CRODET O40)
	17
	POE(20) cetostearyl ether (VOLPO CS20)
	18
20	POE(15) cetostearyl ether (VOLPO CS15)
	19
	POE(20) oleyl ether (VOLPO N20)
	20
	POE(15) oleyl ether (VOLPO N15)
	21
25	POE(45) hydrogenated castor oil (CRODURET 40 or CREMOPHOR RH40)
	22
	POE(35) castor oil (ETOCAS 35)
	23
30	POE(15) glyceryl monolaurate (GLYCEROL L15)

EXAMPLE 24

35

Diclofenac Acid Capsules - Size 1

Capsules of 25 mg active ingredient per capsule are prepared, following generally the procedure of Example 1 but using Size 1 capsules, using the following proportions:

40

45

mg per capsule

Diclofenac acid	25
POE(15) cetostearyl ether (VOLPO CS15)	425
	<hr/>
Total	450

50

55

EXAMPLES 25 TO 27

The procedure of Example 24 was repeated except that 425 mg capsule of the surfactant shown below was used.

<u>Example No</u>	<u>Surfactant</u>
25	POE(20) oleyl ether (VOLPO N20)
26	POE(15) oleyl ether (VOLPO N15)
27	POE(24) monostearate (CRODET S24)

EXAMPLE 28Diclofenac Acid Capsules - Size 0

Capsules of 25 mg active ingredient per capsule are prepared, following generally the procedure of Example 24 but using Size 0 capsules, using the following proportions:

	<u>mg per capsule</u>
Diclofenac acid	25
POE(24) monostearate (CRODET S24)	585
Total	<hr/> 610

EXAMPLES 29 TO 35

The procedure of Example 28 was repeated except that 585 mg. capsule of the surfactant shown below was used.

5

<u>Example No</u>	<u>Surfactant</u>
10 29	POE(40) monostearate (CRODET S40)
30	POE(20) sorbitan monooleate (CRILLET 4)
15 31	POE(20) sorbitan monoisostearate (CRILLET 6)
32	POE(40) hydrogenated castor oil (CRODURET 40 or CREMOPHOR RH40)
20 33	POE(35) castor oil (ETOCAS 35 or CREMOPHOR EL)
34	POE(15) glyceryl monolaurate (GLYCEROX L15)
25 35	POE(20) cetostearyl ether (VOLPO CS20)

30 EXAMPLE 36Piroxicam capsules - Size 1

Following the general procedure of Example 1, except that Size 1 capsules were used, the following capsules were made up.

35

40

mg per capsule

45

Piroxicam	10
POE(20) sorbitan monooleate (CRILLET 4)	440

50

Total	450
-------	-----

55

EXAMPLES 37 TO 44

The procedure of Example 36 was repeated, except that 440 mg capsule of the surfactant shown below was used.

5

<u>Example No</u>	<u>Surfactant</u>
37	POE(20) sorbitan monoisostearate (CRILLET 6)
38	POE(20) cetostearyl ether (VOLPO CS20)
39	POE(15) cetostearyl ether (VOLPO CS15)
40	POE(20) oleyl ether (VOLPO N20)
41	POE(15) oleyl ether (VOLPO N15)
42	POE(40) hydrogenated castor oil (CREMOPHOR RH40)
43	POE(35) castor oil (ETOCAS 35)
44	POE(15) glyceryl monolaurate (GLYCEROX L15)

EXAMPLE 45

30

Ketoprofen Capsules - Size 1

Capsules of 50 mg active ingredient per capsule are prepared in Size 1 gelatin capsules following the general method of Example 1 and using the following proportions:

35

40

	<u>mg per capsule</u>
Ketoprofen	50
POE(20) sorbitan monooleate (CRILLET 4)	400
Total	<hr/> 450

50

55

EXAMPLES 46 TO 51

The procedure of Example 45 was repeated, except that 400 mg capsule of the surfactant shown below was used.

5

<u>Example No</u>	<u>Surfactant</u>
46	POE(20) sorbitan monoisostearate (CRILLET 6)
47	POE(40) monostearate (CRODET S40)
48	POE(24) monostearate (CRODET S24)
49	POE(45) hydrogenated castor oil (CRODURET 40)
50	POE(35) castor oil (ETOCAS 35 or CREMOPHOR EL)
51	POE(24) monolaurate (CRODET L24)

25

EXAMPLE 52Ketoprofen Capsules - Size 2

30

The procedure of Example 45 was repeated, except that Size 2 capsules were used and the ingredients were as follows:

35

	<u>mg per capsule</u>
Ketoprofen	50
POE(20) cetostearyl ether (VOLPO CS20)	285
	<hr/>
Total	335

45

50

55

EXAMPLES 53 TO 58

The procedure of Example 36 was repeated, except that 285 mg capsule of the surfactant shown below was used:

5

<u>Example No</u>	<u>Surfactant</u>
53	POE(15) cetostearyl ether (VOLPO CS15)
54	POE(20) oleyl ether (VOLPO N20)
55	POE(15) oleyl ether (VOLPO N15)
56	POE(40) glyceryl monolaurate (GLYCEROX L40)
57	POE(40) hydrogenated castor oil (CRODURET 40)
58	POE(35) castor oil (ETOCAS 35)

20

It should be noted that if Size 2 capsules formulate satisfactorily then it follows that Size 1 will too.

EXAMPLE 5925 Naproxen Capsules - Size 1

Capsules of 25 mg active ingredient per capsule are prepared in Size 1 gelatin capsules following the general method of Example 1 and using the following proportions:

30

	<u>mg per capsule</u>
Naproxen	25
POE(15) cetostearyl ether (VOLPO CS15)	425
Total	450

40

EXAMPLES 60 TO 62

The procedure of Example 59 was repeated, except that 425 mg capsule of the surfactant shown below was used.

45

- 60 POE(20) cetostearyl ether (VOLPO CS20)
 61 POE(15) oleyl ether (VOLPON15)
 62 POE(20) oleyl ether (VOLPO N20)

50

EXAMPLE 63

55

Flufenamic Acid Capsules - Size 0

Capsules of 50 mg active ingredient per capsule are prepared in Size 0 gelatin capsules following the general method of Example 1 and using the following proportions:

	<u>mg per capsule</u>
Flufenamic Acid	50
POE(24) monolaurate (CRODET L24)	560
	<hr/>
Total	610

EXAMPLES 64 TO 73

The procedure of Example 63 was repeated, except that 560 mg capsule of the surfactant shown below was used:

- 64 POE(24) monostearate (CRODET S24)
- 65 POE(40) monostearate (CRODET S40)
- 66 POE(20) sorbitan monooleate (CRILLET 4)
- 67 POE(20) sorbitan monoisostearate (CRILLET 6)
- 68 POE(4) hydrogenated castor oil (CREMOPHOR RH40)
- 69 POE(15) glyceryl monolaurate (GLYCEROL L15)
- 70 POE(15) cetostearyl ether (VOLPO CS15)
- 71 POE(20) cetostearyl ether (VOLPO CS20)
- 72 POE(15) oleylether (VOLPO N15)
- 73 POE(20) oleylether (VOLPO N20)

EXAMPLE 74Flufenamic Acid Capsules - Size 1

Capsules of 50 mg active ingredient per capsule are prepared in Size 1 gelatin capsules following the general method of Example 1 and using the following proportions:

	<u>mg per capsule</u>
Flufenamic Acid	50
POE(40) hydrogenated castor oil (CREMOPHOR RH40)	400
	<hr/>
Total	450

EXAMPLES 75 TO 77

The procedure of Example 74 was repeated, except that 400 mg capsule of the surfactant shown below was used:

- 75 POE(15) cetostearyl ether (VOLPO CS15)
- 76 POE(20) cetostearyl ether (VOLPO CS20)
- 77 POE(15) oleyl ether (VOLPO N15)

EXAMPLE 78Ibuprofen Capsules - Size 0

Capsules of 50 mg active ingredient per capsule are prepared in Size 0 gelatin capsules following the general method of Example 1 and using the following proportions:

	<u>mg per capsule</u>
Ibuprofen	50
POE(24) monolaurate (CRODET L24)	560
Total	<hr/> 610

EXAMPLES 79 TO 87

The procedure of Example 78 was repeated, except that 560 mg capsule of the surfactant shown below was used:

- 79 POE(24) monostearate (CRODET S24)
- 80 POE(20) sorbitan monooleate (CRILLET 4)
- 81 POE(20) sorbitan monoisostearate (CRILLET 6)
- 82 POE(40) hydrogenated castor oil (CREMOPHOR RH40)
- 83 POE(15) glyceryl monolaurate (GLYCEROX L15)
- 84 POE(15) cetostearyl ether (VOLPO CS15)
- 85 POE(20) cetostearyl ether (VOLPO CS20)
- 86 POE(15) oleyl ether (VOLPO N15)
- 87 POE(15) oleyl ether (VOLPO N20)

EXAMPLE 88

Ibuprofen Capsules - Size 1

Capsules of 50 mg active ingredient per capsule are prepared in Size 1 gelatin capsules following the general method of Example 1 and using the following proportions:

5

	<u>mg per capsule</u>
Ibuprofen	50
POE(24) monolaurate	400
	<hr/>
Total	450

10

15

EXAMPLES 89 TO 94

20

The procedure of Example 88 was repeated, except that 400 mg capsule of the surfactant shown below was used:

25

- 89 POE(20) sorbitan monoisostearate (CRILLET 6)
- 90 POE(40) hydrogenated castor oil (CREMOPHOR RH40)
- 91 POE(15) cetostearyl ether (VOLPO CS15)
- 92 POE(20) cetostearyl ether (VOLPO CS20)
- 93 POE(15) oleyl ether (VOLPON15)
- 94 POE(20) oleyl ether (VOLPO N20)

30

EXAMPLE 95Indomethacin Solution

35

A solution of indomethacin for paediatric or geriatric use may be made according to the following proportions of principal ingredients, the potency being 25 mg per 5 ml, and the dispensed quantity 200 ml:

40

	<u>Quantity per 200 ml</u>
Indomethacin	1.00 g
Surfactant (POE(20) sorbitan monooleate)	20.0 g
Preservative (potassium sorbate)	0.40 g
Sweetener (sodium saccharin)	qs
Citric acid	qs
Flavouring	qs
Water, purified	to 200 ml

50

Approximately half the required water is placed in a suitable container, together with the potassium sorbate (or other suitable preservative), and the sodium saccharin (or other potent sweetener). The solution is stirred and heated continuously to 50-55°C. This forms the aqueous phase.

55

The surfactant (in this example POE (20) sorbitan monooleate eg CRILLET 4 or TWEEN 80) is heated to 50-55°C with continuous stirring in a separate suitable container. The indomethacin is then added and

stirring is continued until 15 minutes after all the active ingredient has dissolved, the temperature being maintained at 50-55°C. This comprises the non-aqueous phase.

The aqueous phase is then added to the non-aqueous phase with continuous stirring. The addition should be fairly rapid. A clear, slightly yellow solution is formed which is then stirred until cool, no further heating being applied after the start of the addition of the aqueous phase to the non-aqueous phase. The solution is then adjusted to give the correct potency by addition of purified water.

pH adjustment is by addition of citric acid until a pH of 3.0-3.5 is reached, the solution being continuously stirred and the citric acid being allowed to completely dissolve before a pH measurement is made. Flavouring is added according to requirements. The solution is then ready for bottling.

EXAMPLES 96 AND 97

Indomethacin solutions are prepared as in Example 95, except that 20g of the following surfactants were used:

- 96 POE(20) sorbitan monoisostearate (CRILLET 6)
97 POE(35) castor oil (CREMOPHOR EL)

EXAMPLE 98

Diclofenac Solution

A solution of diclofenac for paediatric or geriatric use may be made, following the general procedure of Example 95, according to the following proportions of principal ingredients, the potency being 25 mg per 5 ml, and the dispensed quantity 200 ml:

	<u>Quantity per 200ml</u>
Diclofenac Acid	1.00 g
POE(40) hydrogenated castor oil (CREMOPHOR RH40)	27.5 g
Preservative (potassium sorbate)	0.40 g
Sweetener (sodium saccharin)	qs
Citric Acid	qs
Flavouring	qs
Water, purified	to 200 ml

EXAMPLE 99

A diclofenac solution is prepared as in Example 98, except that 27.5 g POE(35) castor oil (CREMOPHOR EL) is used.

EXAMPLE 100

Ketoprofen Solution

A solution of ketoprofen for paediatric or geriatric use may be made following the general procedure of Example 95, according to the following proportions of principal ingredients, the potency being 25 mg per 5 ml, and the dispensed quantity 200 ml:

Quantity per 200 ml

5	Ketoprofen	1.00 g
	Surfactant POE (20) sorbitan	
	monoistearate (CRILLET 6)	10.0 g
10	Preservative (potassium sorbate)	0.40 g
	Sweetener (sodium saccharin)	qs
	Citric acid	qs
15	Flavouring	qs
	Water, purified	to 200 ml

EXAMPLE 101-103

20 A ketoprofen solution is prepared as in Example 100, except that 10g of the following surfactants were used:

- 25 101 POE(40) monostearate (CRODET S40)
 102 POE(20) sorbitan monooleate (CRILLET 4 or TWEEN 80)
 103 POE(40) hydrogenated castor oil (CREMOPHOR RH40)

EXAMPLE 104Flurbiprofen Capsules - Size 1

35 Capsules of 50mg active ingredient per capsule were prepared in Size 1 gelatin capsules following generally the procedure of Example 1 and using the following proportions:

40		<u>mg per capsule</u>
	Flurbiprofen	50
45	POE(40) hydrogenated castor oil	
	(CRODURET 40)	400
50	Total:	<u>450</u>

EXAMPLES 105 TO 109

The procedure of Example 104 was repeated, except that 400mg capsule of the surfactant shown below was used.

<u>Example No.</u>	<u>Surfactant</u>
105	POE(35) castor oil (ETOCAS 35)
106	POE(20) cetostearyl ether (VOLPO CS20)
107	POE(15) cetostearyl ether (VOLPO CS15)
108	POE(20) oleyl ether (VOLPO N20)
109	POE(15) oleyl ether (VOLPO N15)

EXAMPLE 110Flurbiprofen Capsules - Size 0

Following the procedure of Example 104, but using Size 0 capsules, capsules of 50mg active ingredient per capsule were prepared using the following proportions:

	<u>mg per capsule</u>
Flurbiprofen	50
POE(20) sorbitan monooleate (CRILLET 4)	560
Total:	<hr/> 610

EXAMPLE 111 TO 121

The procedure of Example 110 was repeated, except that 560mg capsule of the surfactant shown below was used.

<u>Example No.</u>	<u>Surfactant</u>
111	POE(40) hydrogenated castor oil (CREMOPHOR RH40 or CRODURET 40)
112	POE(35) castor oil (ETOCAS 35 or CREMOPHOR EL)
113	POE(24) monolaurate (CRODET L24)
114	POE(24) monostearate (CRODET S24)
115	POE(20) sorbitan monoisostearate (CRILLET 6)
116	POE(60) hydrogenated castor oil (CREMOPHOR RH60)
117	POE(15) glyceryl monolaurate (GLYCEROX L15)
118	POE(15) cetostearyl ether (VOLPO CS15)
119	POE(20) cetostearyl ether (VOLPO CS20)
120	POE(15) oleyl ether (VOLPO N15)
121	POE(20) oleyl ether (VOLPO N20)

EXAMPLE 122Slow Release Indomethacin Capsules

Capsules of 75mg active ingredient per capsule were prepared using the following proportions:

	<u>mg per capsule</u>
Indomethacin	75
GELUCIRE 46/07	214
POE(24) monostearate [CRODET S24]	321
Total:	<hr/> 610

GELUCIRE 46.07 (by Gattefosse) is a mixture of glycerol and PEG fatty acid esters, with melting point of 43-49°C, HLB of 7, and oral toxicity of LDO > 20g/kg.

The GELUCIRE 46.07 and the POE(24) monostearate were heated, melted and mixed together to 55-60°C and the indomethacin was then added with stirring, the latter being sufficiently vigorous to ensure that the active ingredient was dissolved completely in the mix. The mixture was then filled into hard gelatin

capsules. Size 0.

EXAMPLE 123

The procedure of Example 122 was repeated except that the following ingredients were used in the formulation:

	<u>mg per capsule</u>
Indomethacin	75
GELUCIRE 50/02	214
POE(24) monostearate [CRODET S24]	321
	<hr/>
Total:	610

GELUCIRE 50/02 (by Gattefosse) is a mixture of glycerol and PEG fatty acid esters, with melting point of 48-52°C, HLB of 2, and oral toxicity of LD50 > 18g/kg.

EXAMPLE 124

The procedure of Example 122 was repeated except that the following ingredients were used in this formulation:

	<u>mg per capsule</u>
Indomethacin	75
GELUCIRE 53/10	161
POE(24) monostearate [CRODET S24]	374
	<hr/>
Total:	610

GELUCIRE 53/10 (by Gattefosse) is a mixture of glycerol and fatty acid esters, with melting point of 51-56°C, HLB of 10, and oral toxicity of LDO > 20g/kg.

EXAMPLE 125

The procedure of Example 122 was repeated except that the following ingredients were used in the formulation:

5

10

15

	<u>mg per capsule</u>
Indomethacin	75
GELUCIRE 53/10	214
POE(24) monostearate [CRODET S24]	321
Total:	<u>610</u>

20

EXAMPLE 126

The procedure of Example 122 was repeated except that the following ingredients were used in the formulation:

25

30

35

	<u>mg per capsule</u>
Indomethacin	75
GELUCIRE 53/10	267
POE(24) monostearate [CRODET S24]	268
Total:	<u>610</u>

EXAMPLE 127

40

The procedure of Example 122 was repeated except that the following ingredients were used in the formulation:

45

50

55

	<u>mg per capsule</u>
Indomethacin	75
GELUCIRE 53/10	321
POE(24) monostearate [CRODET S24]	-214
Total:	<u>610</u>

EXAMPLE 128

Capsules from Examples 122 to 127 were assessed for their dissolution rate using USP Apparatus No. 2 (USPXXI) with a paddle speed of 100 rpm, the dissolution medium being 0.2M phosphate buffer pH 7.2 maintained at 37°C.

Aliquots were taken at hourly intervals and the amount of indomethacin dissolved was determined by UV spectrophotometric absorption at 318nm. The results which are the average of three capsules are as follows:

Percentage of Indomethacin dissolved

Time(h)	Example 122	Example 123	Example 124	Example 125	Example 126	Example 127
1	36.0	26.1	31.2	26.4	25.9	19.0
2	59.3	42.6	44.2	37.7	37.4	27.5
3	78.0	54.1	55.2	46.8	44.9	33.7
4	84.5	64.0	66.0	55.1	51.5	39.0
5	90.2	71.9	75.7	63.1	57.5	44.1
6	94.0	78.6	85.1	70.1	63.4	48.7
7	97.0	84.0	91.4	76.4	69.6	52.9
8	98.7	88.6	95.7	81.9	74.7	57.2
9	99.7	90.6	97.7	86.0	79.2	61.8
10	100.0	92.3	98.9	89.8	83.2	66.2
11	100.0	92.8	98.8	92.4	86.3	70.2
12	100.0	92.8	98.4	93.8	88.3	73.6

Claims

1. Micelles containing a non-steroidal anti-inflammatory drug.
2. Micelles as claimed in claim 1, wherein the non-steroidal anti-inflammatory drug is diclofenac, flufenamic acid, flurbiprofen, ibuprofen, indomethacin, ketoprofen, naproxen, phenylbutazone, piroxicam and/or sulindac.
3. A pharmaceutical composition comprising a non-steroidal anti-inflammatory drug and a surfactant, the composition being capable of forming micelles containing the non-steroidal anti-inflammatory drug when administered orally.
4. A composition as claimed in claim 3, wherein the non-steroidal anti-inflammatory drug is diclofenac, flufenamic acid, flurbiprofen, ibuprofen, indomethacin, ketoprofen, naproxen, phenylbutazone, piroxicam and/or sulindac.
5. A composition as claimed in claim 3 or 4, wherein the surfactant is a nonionic surfactant.
6. A composition as claimed in claim 5, wherein the nonionic surfactant is a polyoxyethylated surfactant.
7. A composition as claimed in any one of claims 3 to 6, wherein the surfactant is a polyoxyethylated glycol monoether, a polyoxyethylated fatty acid, a polyoxyethylated sorbitan fatty ester or a polyoxyethylated castor oil.

8. A composition as claimed in any one of claims 3 to 7, wherein the surfactant has an HLB of 10 or above.

9. A composition as claimed in any one of claims 3 to 8, wherein the drug:surfactant weight ratio is in a range of from 1:5.7 to 1:50.

10. A process for the preparation of an anti-inflammatory composition capable of forming non-steroidal anti-inflammatory drug-containing micelles on oral administration to a human or non-human animal, the process comprising admixing a non-steroidal antiinflammatory drug with a surfactant.

11. The use of a non-steroidal anti-inflammatory drug and a surfactant in the preparation of a composition for administering the drug in micellar form.

Claims for the following Contracting States: ES and GR

1. A process for the preparation of an anti-inflammatory composition capable of forming non-steroidal anti-inflammatory drug-containing micelles on oral administration to a human or non-human animal, the process comprising admixing a non-steroidal anti-inflammatory drug with a surfactant.

2. A process as claimed in claim 1, wherein the non-steroidal anti-inflammatory drug is diclofenac, flufenamic acid, flurbiprofen, ibuprofen, indomethacin, ketoprofen, naproxen, phenylbutazone, piroxicam and or sulindac.

3. A process as claimed in claim 1 or 2, wherein the surfactant is a nonionic surfactant.

4. A process as claimed in claim 3, wherein the nonionic surfactant is a polyoxyethylated surfactant.

5. A process as claimed in any one of claims 1 to 4, wherein the surfactant is a polyoxyethylated glycol monoether, a polyoxyethylated fatty acid, a polyoxyethylated sorbitan fatty ester or a polyoxyethylated castor oil.

6. A process as claimed in any one of claims 1 to 5, wherein the surfactant has an HLB of 10 or above.

7. A process as claimed in any one of claims 1 to 6, wherein the drug:surfactant weight ratio is in a range of from 1:5.7 to 1:50.

8. The use of a non-steroidal anti-inflammatory drug and a surfactant in the preparation of a composition for administering the drug in micellar form.

THIS PAGE BLANK (USPTO)

12

EUROPEAN PATENT APPLICATION

21 Application number: 87310931.8

51 Int. Cl.⁴: A 61 K 9/10
A 61 K 47/00

22 Date of filing: 11.12.87

30 Priority: 18.12.86 GB 8630273

43 Date of publication of application:
20.07.88 Bulletin 88/29

84 Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

89 Date of deferred publication of search report:
07.09.88 Bulletin 88/36

71 Applicant: T.I.L. Medical Ltd.
The Old Blue School Lower Square
Isleworth Middlesex TW7 6RL (GB)

72 Inventor: Story, Michael John
Elm Cottage Greaves Lane
Threapwood Near Malpas Cheshire SY14 6AS (GB)

Flynn, Michael John
Hunterscombe Dorking Road
Leatherhead Surrey KT22 8JT (GB)

74 Representative: Sheard, Andrew Gregory et al
Kilburn & Strode 30, John Street
London WC1N 2DD (GB)

54 Micelles containing a non-steroidal antiinflammatory compound.

57 Non-steroidal anti-inflammatory drugs (NSAIDs) including diclofenac, flufenamic acid, flurbiprofen, ibuprofen, indomethacin, ketoprofen, naproxen, phenylbutazone, piroxicam and sulindac are administered in micelles to alleviate their adverse effects on the gastrointestinal tract. The drugs are formulated with surfactants such as polyethoxylated nonionics to give micelle-forming compositions.

EP 0 274 870 A3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 87 31 0931

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
P, X	EP-A-0 223 369 (THE BOOTS CO. PLC) * Page 2, line 30 - page 3, line 1; page 3, lines 32-35; page 4, lines 24-32; page 8, lines 27-33; claim 1 *	1-11	A 61 K 9/10 A 61 K 47/00
X	EP-A-0 178 436 (DOLORGIET GmbH) * Examples 1,6,11,13,14,16,18; claims 1-4 *	1-8,10, 11	
X	EP-A-0 179 583 (MERCK & CO. INC.) * Page 1, lines 1-5; page 2, line 10 - page 3, line 24; page 5, lines 35-37; page 7, lines 13-34; page 10, lines 14-18; claims 1-3 *	1-11	
X	CHEMICAL ABSTRACTS, vol. 90, no. 16, 16th April 1979, page 370, abstract no. 127464t, Columbus, Ohio, US; H. KRASOWSKA: "Solubilities of certain antiinflammatory compounds in nonionic surfactant solutions", & PHARM. IND. 1978, 40(12), 1381-4 * Abstract *	1-11	TECHNICAL FIELDS SEARCHED (Int. Cl.4)
X	CHEMICAL ABSTRACTS, vol. 94, no. 16, 10th April 1981, page 378, abstract no. 127260f, Columbus, Ohio, US; H. KRASOWSKA: "Effect of micellar solubilization on the gastrointestinal absorption of indomethacin in the rat", & INT. J. PHARM. 1980, 7(2), 137-43 * Abstract *	1-11	A 61 K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 16-06-1988	Examiner MUELLNERS W.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	



European Patent
Office

EUROPEAN SEARCH REPORT

Page 2

Application Number

EP 87 31 0931

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
X	CHEMICAL ABSTRACTS, vol. 94, no. 8, 23rd February 1981, page 358, abstract no. 52772p, Columbus, Ohio, US; A.H. GHANEM et al.: "Solubilization of flufenamic acid", & PHARM. IND. 1980, 42(8), 854-6 * Abstract * -----	1-11	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 16-06-1988	Examiner MUELLNERS W.
CATEGORY OF CITED DOCUMENTS			
<p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons</p> <p>& : member of the same patent family, corresponding document</p>			

THIS PAGE BLANK (USPTO)